CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 102, 2009

A

Abel, C. A., 629
Aggamy, E., 809
Aghdam, H. R., 233
Aguayo, I., 426
Akbar, W., 96
Akman, O., 467
Alarie, Y., 797
Albert, P. J., 717
Alexandrakis, V., 835
Allahyari, H., 450
Al-Lawati, H., 881
Aluja, M., 560
Al-Wahaibi, A. K., 241, 249
Ammar, E.-D., 889
Antonini, G., 998
Armbruster, P., 1135
Armstrong, J. E., 492
Armstrong, J. S., 196
Ascher, J. S., 275
Aslin, R., 426
Atwa, A., 809
Audisio, P., 998
Augot, D., 347

B

Babba, H., 347
Backus, E. A., 847
Baez, I., 445
Bailowitz, R. A., 261
Bardunias, P., 158, 906
Barrera, J. F., 166
Barta, R. J., 925
Barták, M., 396
Bauce, E., 717
Bell, A. J., 913
Bennett, G. W., 1029
Berchi, S., 347
Berestecky, J. M., 498
Berlocher, S. H., 381
Bernhardt, J. L., 638
Bienefeld, K., 881
Bischoff, I., 60
Blanco, C. A., 629
Blochtein, B., 137
Boggs, C. L., 1105
Boina, D. R., 523
Boudabous, R., 347
Bounamous, A., 347
Boyd, D. W., 386
Brailovsky, H., 959
Braman, S. K., 386
Brinton, J., 467
Broufas, G. D., 70, 437
Brown, B. V., 937
Brown, J. K., 650
Brust, M. L., 462
Bu, C.-P., 295
Buczkowski, G., 1029
Bundy, C. S., 1068

C

Cabrera Walsh, G., 603
Cardinal-Aucoin, M., 717
Carlton, C., 96, 406
Carter, M. E., 895
Cassill, D. L., 713
Castrejón-Gómez, V. R., 310
Chapman, P. S., 1013
Chen, X.-L., 674
Chen, X.-X., 532
Chen, Z., 386
Chiel, E., 413
Childers, C. C., 617
Chiu-Alvarado, P., 166
Christiansen, K., 1037
Cicero, J. M., 650
Cline, A. R., 998
Cohnstaedt, L., 144
Coleman, R. J., 196
Collier, G. E., 492
Compton, S. G., 556
Conle, O. V., 210
Cordero-Rivera, A., 694
Córdoba-Aguilar, A., 694
Cossé, A. A., 560
Couloux, A., 347

D

Daane, K. M., 727
D'adamo, P., 1109
De Biase, A., 998
Delaney, D. A., 666
Delebecque, C. J., 925
Depaquit, J., 347
Desneux, N., 925
Díaz, N. B., 603
Díez, J. J., 317
Dossey, A. T., 210
Downie, D. A., 1126
Duncan, R. E., 112
Durham, G., 426

E

Eastop, V. F., 914
Eaton, T. D., 386
Eberhard, S. H., 1105
El-Aziz, G.M.A., 809
El-Rahman, S.M.A., 809
Equihua-Martinez, A., 597
Esquivel, J. F., 303

F

Fadamiro, H. Y., 617
Fathipour, Y., 233
Favret, C., 970
Ferté, H., 347
Fialho, M.d.C.Q., 119
Franklin, M. T., 151
Freitas, A.V.L., 1151
Fujisaki, K., 76
Furgol, M., 1037
Furniss, M. M., 597

G

Gaddi, A. L., 603
Gallardo, F. E., 603
Gao, T., 588
Gariepy, T. D., 925
Gemeno, C., 317
Gilbert, L. E., 937
Giroux, M., 567
Gonçalves dos Santos, C., 137
Grace, J. K., 498
Gray, D. A., 367
Groot, A. T., 629
Gruwell, M. E., 1050
Gülgönül, E. E., 288

H

Hajek, A. E., 913
Hall, D. G., 476, 547
Hamilton, G. C., 608
Hamud, S. M., 426
Hanks, L. M., 381
Hannam, J. J., 913
Hanula, J. L., 508
Harrison, R. G., 895
Haruyama, N., 747
Harvey, J. A., 172
Hayes, J. L., 982
He, J.-H., 532
Hegazi, E., 809
Heimpel, G. E., 925
Hennemann, F. H., 210
Henry, C. S., 747
Hermann, S. L., 523
Hikl, A. L., 1105
Hoback, W. W., 462
Hoddle, M., 835
Hoelmer, K. A., 925
Holmes, W. J., 847
Hong, F., 295
Hong, X.-Y., 674
Hori, S., 826
Houseman, R. M., 868
Huang, F., 532

Hunter, M. S., 413
Husband, D. O., 1062
Husband, R. W., 1062
Husseneder, C., 498

I

Ibeas, F., 317
Inbar, M., 413
Isaacs, R., 275
Ivanović, A., 1

J

Jacas, J.-A., 112
Jannot, J. E., 467
Janzen, D. H., 766
Jenkins, T. M., 386
Johnson, M. W., 727
Johnson, P. J., 735
Jones, J. C., 181
Jones, S. C., 889
Jouet, D., 347
Juliano, S. A., 467

K

Kang, J., 326
Kanzaki, N., 684, 759
Karamaouna, F., 1145
Katakura, H., 826
Kato, T., 826
Kaufman, L. V., 104
Kavallieratos, N. G., 1
Keena, M. A., 539
Kennedy, J. H., 1165
Khafagi, W. E., 809
Kobayashi, N., 826
Kocot, K., 467
Kongchuensin, M., 360
Konstantopoulou, M., 809, 1145
Kontodimas, D. C., 233, 1145
Koveos, D. S., 70, 437
Kral, K., 339
Krenn, H. W., 1105
Krupke, C. H., 326
Kubík, Š., 396

L

Labrzycka, A., 12
Lamanna, F., 998
Lauzière, I., 642
Lauzon, C. R., 85
Lebrun, E. G., 937
Lee, C.-F., 48, 785
Lee, C.-Y., 1077, 1091
Lee, J. C., 426
Lee, S.-H., 1158
Legaspi, B. C., Jr, 445
Legaspi, J. C., 445
Li, H.-F., 333, 684, 759
Li, S., 1029
Li, T., 674
Li, Z.-X., 39
Liang, A.-P., 739
Liebherr, J. K., 913
Liu, C., 20
Lozada, M., 1109

M

Maginnis, T. L., 1099
Makarkin, V. N., 964
Mancini, E., 998
Mankin, R. W., 547
Martinou, A. F., 1145
Mason, A., 736
McCombs, S. D., 85
McPherson, J. E., 1068
Medeiros-Santana, L., 819
Megiolaro, F. L., 137
Meixner, M. D., 666
Mendenhall, M., 181
Messina, F. J., 181
Metz, M. A., 766
Michat, M. C., 797
Milonas, P. G., 1145
Mirmohammadi, S., 450
Misof, B., 60
Mitchell, R. F., 381
Mochizuki, A., 747
Moltzan, B. D., 426
Morales-Ramos, J., 629
Morse, J. G., 241, 249
Moser, S. E., 419
Mozes-Daube, N., 413
Muller, A., 181
Munson, A. S., 426
Munstermann, L. E., 144
Myers, J. H., 151

N

Nakahira, K., 747
Nakamatsu, Y., 172
Negrón, J. F., 426
Nematollahi, M. R., 450
Neoh, K.-B., 1091
Nesbitt, M., 617
Neves, C. A., 119
Nie, J., 739
Nielsen, A. L., 608
Nielsen, C., 913
Nishida, T., 201
Nomura, M., 747
Normark, B. B., 1050

O

Obrycki, J. J., 419
O'Donnell, B. C., 128
O'Donnell, D., 1135
Ohta, Y., 826
Oliveira, P. S., 1151
Onagbola, E. O., 523
Opp, S., 727
Othman, A. S., 1077
Ouyang, X.-H., 295
Ozaki, K., 456

P

Pajares, J. A., 317
Pappas, M. L., 70, 437
Parrella, M. P., 484
Peabody, N. C., 85
Peña, J. E., 112
Peng, Y. Q., 556
Peredo, L. C., 959
Peterson, T., 426
Pettis, G. V., 386
Philpott, D. E., 381
Pinzon, O. P., 868
Plowes, R. M., 937
Pomorski, R. J., 1037
Potter, S. E., 85

Q

Qiu, L.-F., 39

R

Radjabi, G., 233
Ragenovich, I. R., 426
Raimundo, R.L.G., 1151
Rakhshani, E., 1
Ramalho, F. S., 119
Rao, A., 702
Raptopoulos, D., 809
Reagan, T. E., 96
Reardon, B. J., 847
Rector, B. G., 12
Redmond, C. R., 1099
Ren, D., 20, 588, 964
Rezapanah, M., 233
Richardson, M. L., 737
Ridland, P. M., 914
Ritland, C. E., 151
Robacker, D. C., 560
Roberts, P. D., 650
Rojas, J. C., 166, 310
Rojas, M. G., 629
Ruiz, E. A., 982

S

Saboori, A., 450
Sacchetti, P., 560
Sahara, K., 360
Saito, Y., 360
Sak, O., 288
Sakagami, T., 360
Sánchez, V., 539
Sano, M., 456
Savage, J., 354
Scharf, M. E., 1029
Schiff, N. M., 666
Schreiber, F., 847
Schröder, S., 60
Schwinghammer, M., 1029
Serrano-Meneses, M. A., 694
Serrão, J. E., 119, 137
Sétamou, M., 196, 642
Seybold, S. J., 426
Sheets, H. D., 1013

Sheppard, W. S., 666
Shi, M., 532
Shields, V.D.C., 1116
Shih, C., 20, 588
Shimizu, K., 76
Showeil, S., 809
Singh, D., 713
Skoracka, A., 12
Smith, M. T., 895
Solis, M. A., 766
Stanisavljević, L. Ž., 1
Stansly, P. A., 650
Starý, P., 1, 925
Stelinski, L. L., 476, 523
Stevens, L. E., 261
Su, N.-Y., 158, 333, 684, 759, 906, 1158
Sun, J., 508
Sutherland, A. M., 484
Suzuki, S., 826

T

Tabuchi, K., 456
Tanaka, T., 172
Tawfik, H., 809
Thomas, J. C., 189
Tomanović, Ž., 1
Trager, J. C., 911
Trizzino, M., 998
Tsitsipis, I., 835
Tuell, J. K., 275

U

Uçkan, F., 288
Unruh, T. R., 1013
U¨mani-Brown, S., 144

V

Valenzuela, I., 914
Varikou, K., 835
Veuille, M., 347
Víctor, J., 982
Vinson, S. B., 702

W

Walker, G. P., 847
Walker, T. J., 367
Wang, C., 1029
Wang, H.-L., 295
Wang, X.-G., 727
Watanabe, H., 517
Watrous, L. E., 406
Weeks, A. R., 914
Weissman, D. B., 367
Wenninger, E. J., 476, 547
Wheeler, T. A., 567
White, J. A., 413
Williams, J. G., 1126
Witcosky, J. J., 426
Wright, M. G., 104
Wu, J., 1050

X

Xi, G.-S., 295
Xiao, Y., 617

Y

Yamasaki, A., 76
Yamauchi, S., 826
Yang, D. R., 556
Yang, Q., 964
Yang, R.-L., 906
Yang, Z.-Q., 39
Yano, E., 517
Yao, Y.-X., 39
Ye, G.-Y., 532
Ye, W., 684
Ye, Y., 889
Yeap, B.-K., 1077
Yee, W. L., 1013
Yoshimoto, J., 201
Young, O. P., 842

Z

Zanuncio, J. C., 119
Zchori-Fein, E., 413
Zhan, G.-J., 295
Zhang, F. P., 556
Zhang, Y., 508
Zhou, X., 1029
Žikić, V., 1
Zucoloto, F. S., 819
Zúñiga, G., 982

CUMULATIVE KEY WORD INDEX
Annals of the Entomological Society of America
Volume 102, 2009

A

abdominal gills,
  Ephoron leukon, 128
  mandibular tusks, 128
  mayfly ontogeny, 128
  nymphal development, 128
Acaralox,
  Acari, 12
  Aculodes, 12
  Eriophyidae, 12
  Poaceae, 12
Acari, 12, 1062
  Acaralox, 12
  Aculodes, 12
  Carabidae, 1062
  Eriophyidae, 12
  Poaceae, 12
  Podapolipidae, 1062
  taxonomy, 1062
acoustic communication,
  Asian citrus psyllid, 547
  citrus greening disease, 547
  huanglongbing, 547
acridids,
  Caelifera, 339
  catantopids, 339
  Orthoptera, 339
  pygmy grasshoppers, 339
active galleries,
  mass trapping, 809
  olive yield, 809
  sticky UV-light–pheromone trap, 809
  Zeuzera pyrina, 809
activity schedule,
  ant behavior, 1151
  diet, 1151
  ground-dwelling ants, 1151
Aculodes,
  Acaralox, 12
  Acari, 12
  Eriophyidae, 12
  Poaceae, 12
adaptation,
  artificial selection, 181
  experimental evolution, 181
  host shift, 181
  oviposition, 181
Adelges tsugae,
  allozymes, 539
  genetic diversity, 539
  parthenogenic, 539
adult body weights,
  egg incubation, 445
  egg production, 445
  Opuntia ficus-indica, 445
adventive species,
  life cycle, 104
  Pipturus spp., 104
  purposely introduced species, 104
Aedes albopictus, 144, 1135
  invasive species, 144, 1135
  life history, 1135
  mitochondrial DNA gene, 144
  nicotinamide adenine dinucleotide dehydrogenase subunit 5, 144
  population genetics, 144
  population structure, 1135
aestivation,
  extended development, 456
  galling aphid, 456
  life history evolution, 456
  Pinaceae, 456
aggression,
  coexistence, 201
  competitor avoidance, 201
  dominance hierarchy, 201
  interference competition, 201
alates,
  oocyte, 889
  reproduction, 889
  sperm energy source, 889
  spermatheca, 889
Aleocharinae,
  12s, 189
  16s, 189
  rove beetles, 189
  Staphylinidae, 189
alimentary canal,
  filter chamber, 650
  huanglongbing, 650
  Liberibacter, 650
  Psyllidae, 650
allozyme, 381, 539
  Adelges tsugae, 539
  genetic diversity, 539
  mitochondrial DNA, 381
  morphology, 381
  parthenogenic, 539
  phytophagous insect, 381
  systematics, 381
Altica litigata,
  flea beetle, 386
  herbivore, 386
  internal transcribed spacer, 386
  mitochondrial DNA, 386
amplified fragment length polymorphism,
  Bt resistance, 151
  genetic variation, 151
  population structure, 151
  Trichoplusia ni, 151
Anastrepha serpentina,
  antennal morphology, 310
  fruit flies, 310
  olfaction, 310
  sensilla, 310
Anisomorpha buprestoides,
  color forms, 210
  ecology, 210
  host plants, 210
  intraspecific variation, 210
Annonaceae,
  Costa Rica, 492
  Drosophilidae, 492
  florivory, 492
  saproflorivory, 492
Anoplophora glabripennis,
  Asia, 895
  Asian longhorned beetle, 895
  microsatellites, 895
  mitochondrial DNA, 895
ant behavior,
  activity schedule, 1151
  diet, 1151
  ground-dwelling ants, 1151
antennal morphology, 310, 523
  Anastrepha serpentina, 310
  citrus greening, 523
  electron microscopy, 523
  fruit flies, 310
  huanglongbing, 523
  olfaction, 310
  sensilla, 310
antipredator strategy,
  behavior, 517
  crypsis, 517
  mantid, 517
  wind, 517
Apis mellifera,
  commercial populations, 666
  genetic diversity, 666
  microsatellites, 666
  mitochondrial DNA, 666
apple maggot,
  canonical variates analysis, 1013
  geometric morphometrics, 1013
  ovipositor length, 1013

Argopistes tsekooni,
biological control, 508
life history, 508
Ligustrum sinense, 508
artificial selection,
adaptation, 181
experimental evolution, 181
host shift, 181
oviposition, 181
Asia, 895, 1077
Anoplophora glabripennis, 895
Asian longhorned beetle, 895
Coptotermes, 1077
microsatellites, 895
mitochondrial DNA, 895, 1077
phylogenetics, 1077
subterranean termites, 1077
Asian citrus psyllid, 476, 547
acoustic communication, 547
body color, 476
citrus greening disease, 547
fecundity, 476
huanglongbing, 547
pigment, 476
reproductive output, 476
Asian longhorned beetle,
Anoplophora glabripennis, 895
Asia, 895
microsatellites, 895
mitochondrial DNA, 895
Asteraceae,
Meligethinae, 998
Nitidulidae, 998
systematics, 998
Tarchonantheae, 998
Aubehydrini,
chaetotaxy, 797
diving beetles, 797
Dytiscidae, 797
phylogeny, 797
Auchenorrhyncha,
development, 642
life cycle, 642
reproduction, 642
Xylella fastidiosa, 642

B

Bactrocera (*Dacus*) *oleae*,
egg production, 70
longevity, 70
ovarian maturation, 70
relative humidity, 70
behavior, 137, 517, 702
antipredator strategy, 517
crypsis, 517
Forelius, 702
gland plasticity, 137
life stages, 137
mantid, 517
Monomorium, 702
Pheidole, 702
propolis, 137
stingless bees, 137
Tetramorium, 702
wind, 517
Binodoxys communis,
Binodoxys koreanus, 925
cryptic species, 925
parasitoids, 925
soybean aphid, 925
Binodoxys koreanus,
Binodoxys communis, 925
cryptic species, 925
parasitoids, 925
soybean aphid, 925
biocontrol,
egg parasitoid, 39
Halyomorpha halys, 39
new species, 39
Trissolcus halyomorphae, 39
biodiversity,
reproductive isolation, 360
social spider mites, 360
Tetranychidae, 360
biogeography, 20, 261, 684
12S rRNA, 684
16S rRNA, 684
elevation, 261
Grand Canyon, 261
Jiulongshan Formation, 20
Middle Jurassic, 20
mitochondrial COII, 684
molecular phylogeny, 684
Odonata, 261
palaeoentomology, 20
Pelecinidae, 20
species richness, 261
biological control, 96, 112, 241, 484, 508
Argopistes tsekooni, 508
Diaprepes abbreviatus, 112
Diomus terminatus, 96
egg age, 241
egg parasitoid, 112, 241
egg structure, 241
leafhoppers, 241
life cycle, 96
life history, 508
Ligustrum sinense, 508
Melanaphis sacchari, 96
morphology, 96
Pachnaeus litus, 112
powdery mildew, 484
Psyllobora vigintimaculata, 484
biological invasions,
cognition, 1109
memory, 1109
biotypes,
Ephedrus persicae, 1
geometric morphometrics, 1
variability, 1
body color, 76, 476
Asian citrus psyllid, 476
fecundity, 476
Helicoverpa armigera, 76
host plant, 76
larval diet, 76
pigment, 476
polymorphism, 76
reproductive output, 476
Brevicoryne brassicae,
demography, 450
jackknife, 450
life table, 450
oilseed rape, 450
Bt resistance,
amplified fragment length polymorphism, 151
genetic variation, 151
population structure, 151
Trichoplusia ni, 151

C

Caelifera,
acridids, 339
catantopids, 339
Orthoptera, 339
pygmy grasshoppers, 339
Callainireinwardtia woodi,
Muscidae, 354
Neotropical, 354
new genus, 354
Reinwardtiini, 354
canonical variates analysis,
apple maggot, 1013
geometric morphometrics, 1013
ovipositor length, 1013
Capparaceae,
Costa Rica, 766
larval morphology, 766
Pyraloidea, 766
systematics, 766
Carabidae,
Acari, 1062
Podapolipidae, 1062
taxonomy, 1062
caste,
development, 1091
Macrotermes carbonarius, 1091
Macrotermes gilvus, 1091
sexual dimorphism, 1091
catantopids,
acridids, 339
Caelifera, 339
Orthoptera, 339
pygmy grasshoppers, 339
cave,
Collembola, 1037
new species, 1037
Onychiurus, 1037
Cerambycidae,
contact pheromone, 317
cuticular hydrocarbon, 317
mating behavior, 317

Monochamus galloprovincialis, 317
chaetotaxy,
Aubehydrini, 797
diving beetles, 797
Dytiscidae, 797
phylogeny, 797
China, 588, 964
Daohugou, 964
Jiulongshan Formation, 588
Jurassic, 964
Kalligrammatidae, 964
Middle Jurassic, 588
Neuroptera, 964
new genera, 588
Xyelotomidae, 588
Choristoneura fumiferana,
condensed tannins, 717
hydrolysable tannins, 717
insect herbivore, 717
spruce budworm, 717
Cicadomorpha,
drawing, 739
Machaerotinae, 739
new species, 739
taxonomy, 739
Cicindela,
flooding, 462
hypoxia, 462
immersion, 462
tiger beetles, 462
citrus,
developmental time, 835
Pezothrips kellyanus, 835
temperature thresholds, 835
thermal requirements, 835
citrus greening,
antennal morphology, 523
electron microscopy, 523
huanglongbing, 523
citrus greening disease,
acoustic communication, 547
Asian citrus psyllid, 547
huanglongbing, 547
Coccinella septempunctata,
Coleomegilla maculata, 419
Harmonia axyridis, 419
interspecific interactions, 419
introduced species, 419
codling moth,
fertility, 233
life table parameters, 233
reproduction, 233
Weibull distribution, 233
coexistence,
aggression, 201
competitor avoidance, 201
dominance hierarchy, 201
interference competition, 201
cognition,
biological invasions, 1109
memory, 1109
COI,
dung beetle, 826
flightless, 826
molecular phylogeny, 826
Coleomegilla maculata,
Coccinella septempunctata, 419
Harmonia axyridis, 419
interspecific interactions, 419
introduced species, 419
Collembola,
cave, 1037
new species, 1037
Onychiurus, 1037
color changes,
development, 638
Oebalus pugnax, 638
rice stink bug eggs, 638
temperature, 638
color forms,
Anisomorpha buprestoides, 210
ecology, 210
host plants, 210
intraspecific variation, 210
commercial populations,
Apis mellifera, 666
genetic diversity, 666
microsatellites, 666
mitochondrial DNA, 666
compensatory growth,
early embryonic mortality, 881
embryogenesis, 881
maternal effects, 881
queen age, 881
competition,
idiobiont, 172
occupation, 172
ovipositional resource, 172
competitive displacement,
flight behavior, 426
Scolytidae, 426
Scolytus multistriatus, 426
Scolytus schevyrewi, 426
competitor avoidance,
aggression, 201
coexistence, 201
dominance hierarchy, 201
interference competition, 201
condensed tannins,
Choristoneura fumiferana, 717
hydrolysable tannins, 717
insect herbivore, 717
spruce budworm, 717
conservation,
cycad, 1126
cytochrome oxidase I, 1126
elongation factor 1a genes, 1126
endangered species, 1126
contact pheromone,
Cerambycidae, 317
cuticular hydrocarbon, 317
mating behavior, 317
Monochamus galloprovincialis, 317
Coptotermes formosanus, 333, 1158
excavating behavior, 333
individual distance, 1158
individual volume, 1158
mouthparts, 333
Reticulitermes flavipes, 1158
self-organized behavior, 1158
tunnel architecture, 333
Coptotermes, 498, 1077
Asia, 1077
gut symbionts, 498
insect gut, 498
Isoptera, 498
mitochondrial DNA, 1077
phylogenetics, 1077
Rhinotermitidae, 498
subterranean termites, 1077
copulation duration,
mating system, 694
Odonata, 694
Corimelaena incognita,
descriptions, 1068
laboratory rearing, 1068
life history, 1068
Thyreocoridae, 1068
Costa Rica, 492, 766
Annonaceae, 492
Capparaceae, 766
Drosophilidae, 492
florivory, 492
larval morphology, 766
Pyraloidea, 766
saproflorivory, 492
systematics, 766
cotton,
Creontiades signatus, 196
okra-leaf, 196
oviposition, 196
trichomes, 196
Creontiades signatus,
cotton, 196
okra-leaf, 196
oviposition, 196
trichomes, 196
crypsis,
antipredator strategy, 517
behavior, 517
mantid, 517
wind, 517
cryptic,
hybridization, 914
karyotype, 914
mitochondrial DNA, 914
nuclear DNA, 914
cryptic species, 367, 747, 925
Binodoxys communis, 925
Binodoxys koreanus, 925
Gryllus, 367
introduced species, 367
mating signal, 747

parasitoids, 925
sister species, 367
song species, 747
soybean aphid, 925
substrate vibration, 747
systematics, 747
cucurbit domestication,
ethnobiology, 60
geometric morphometrics, 60
range expansion, 60
squash bee, 60
cuticular hydrocarbon,
Cerambycidae, 317
contact pheromone, 317
mating behavior, 317
Monochamus galloprovincialis, 317
cycad,
conservation, 1126
cytochrome oxidase I, 1126
elongation factor 1a genes, 1126
endangered species, 1126
cytochrome oxidase I,
conservation, 1126
cycad, 1126
elongation factor 1a genes, 1126
endangered species, 1126

D

Daohugou,
China, 964
Jurassic, 964
Kalligrammatidae, 964
Neuroptera, 964
degree-days,
developmental threshold, 249
glassy-winged sharpshooter, 249
host plants, 249
demographic parameters,
development, 437
Dichochrysa prasina, 437
Myzus persicae, 437
prey availability, 437
demography,
Brevicoryne brassicae, 450
jackknife, 450
life table, 450
oilseed rape, 450
Dendroctonus pseudotsugae,
mitochondrial DNA, 982
Pseudotsuga, 982
subspecies, 982
descriptions,
Corimelaena incognita, 1068
laboratory rearing, 1068
life history, 1068
Thyreocoridae, 1068
development, 437, 467, 638, 642, 1091
Auchenorrhyncha, 642
caste, 1091
color changes, 638
demographic parameters, 437
Dichochrysa prasina, 437
growth rate, 467
hatching size, 467
interpopulation variation, 467
life cycle, 642
Macrotermes carbonarius, 1091
Macrotermes gilvus, 1091
Myzus persicae, 437
Oebalus pugnax, 638
prey availability, 437
reproduction, 642
rice stink bug eggs, 638
sexual dimorphism, 1091
south Florida, 467
temperature, 638
Xylella fastidiosa, 642
developmental threshold,
degree-days, 249
glassy-winged sharpshooter, 249
host plants, 249
developmental time,
citrus, 835
Pezothrips kellyanus, 835
temperature thresholds, 835
thermal requirements, 835
Diabrotica virgifera virgifera,
mate choice, 326
mating behavior, 326
reproductive behavior, 326
Diadegma semiclausum,
Ichneumonidae, 532
immature stages, 532
Plutella xylostella, 532
preimaginal development, 532
Diaprepes abbreviatus,
biological control, 112
egg parasitoid, 112
Pachnaeus litus, 112
Dichochrysa prasina,
demographic parameters, 437
development, 437
Myzus persicae, 437
prey availability, 437
diet,
activity schedule, 1151
ant behavior, 1151
ground-dwelling ants, 1151
dihydropyrazine,
2,5-dimethylpyrazine, 560
trimethylpyrazine, 560
2,5-dimethylpyrazine,
dihydropyrazine, 560
trimethylpyrazine, 560
Diomus terminatus,
biological control, 96
life cycle, 96
Melanaphis sacchari, 96
morphology, 96
Diptera, 396, 567
Empididae, 396
Israel, 396
Neobellieria, 567
new species, 396
Rhamphomyia, 396
Sarcophaga, 567
systematics, 567
Tolucamyia, 567
discriminant functions,
DNA barcodes, 970
Mindarus abietinus, 970
diving beetles,
Aubehydrini, 797
chaetotaxy, 797
Dytiscidae, 797
phylogeny, 797
DNA barcodes, 347, 970
discriminant functions, 970
Mindarus abietinus, 970
mitochondrial DNA, 347
Phlebotominae, 347
Phlebotomus chabaudi, 347
Phlebotomus riouxi, 347
DNA sequence,
intraspecific variation, 868
Missouri, 868
Reticulitermes, 868
subterranean termites, 868
dominance hierarchy,
aggression, 201
coexistence, 201
competitor avoidance, 201
interference competition, 201
drawing,
Cicadomorpha, 739
Machaerotinae, 739
new species, 739
taxonomy, 739
Drosophilidae,
Annonaceae, 492
Costa Rica, 492
florivory, 492
saproflorivory, 492
dung beetle,
COI, 826
flightless, 826
molecular phylogeny, 826
dust/frass,
ectoparasitoid, 166
host searching, 166
semiochemicals, 166
Dytiscidae,
Aubehydrini, 797
chaetotaxy, 797
diving beetles, 797
phylogeny, 797

E
early embryonic mortality,
compensatory growth, 881
embryogenesis, 881
maternal effects, 881
queen age, 881
ecology,
Anisomorpha buprestoides, 210
color forms, 210
host plants, 210
intraspecific variation, 210
ectoparasitoid,
dust/frass, 166
host searching, 166
semiochemicals, 166
Egeria densa,
Ephydridae, 603
Eucoilinae, 603
Hydrellia, 603
Hydrelliaeucoila egeria, 603
egg age,
biological control, 241
egg parasitoids, 241
egg structure, 241
leafhoppers, 241
egg incubation,
adult body weights, 445
egg production, 445
Opuntia ficus-indica, 445
egg parasitoid, 39, 112, 241, 1145
biocontrol, 39
biological control, 112, 241
Diaprepes abbreviatus, 112
egg age, 241
egg structure, 241
Halyomorpha halys, 39
infochemicals, 1145
leafhoppers, 241
new species, 39
Pachnaeus litus, 112
Prays oleae, 1145
Trissolcus halyomorphae, 39
Y-tube olfactometer, 1145
egg production, 70, 445
adult body weights, 445
Bactrocera (*Dacus*) *oleae*, 70
egg incubation, 445
longevity, 70
Opuntia ficus-indica, 445
ovarian maturation, 70
relative humidity, 70
egg structure,
biological control, 241
egg age, 241
egg parasitoids, 241
leafhoppers, 241
electrical penetration graph,
Homalodisca spp., 847
Pierce's disease, 847
probing, 847
stylet penetration, 847
electron microscopy,
antennal morphology, 523
citrus greening, 523
huanglongbing, 523
elevation,
biogeography, 261
Grand Canyon, 261
Odonata, 261
species richness, 261
elongation factor 1a genes,
conservation, 1126
cycad, 1126
cytochrome oxidase I, 1126
endangered species, 1126
embryogenesis,
compensatory growth, 881
early embryonic mortality, 881
maternal effects, 881
queen age, 881
Empididae,
Diptera, 396
Israel, 396
new species, 396
Rhamphomyia, 396
Encarsia,
horizontal transfer, 1050
parthenogenesis, 1050
secondary endosymbiont, 1050
endangered species,
conservation, 1126
cycad, 1126
cytochrome oxidase I, 1126
elongation factor 1a genes, 1126
Ephedrus persicae,
biotypes, 1
geometric morphometrics, 1
variability, 1
Ephoron leukon,
abdominal gills, 128
mandibular tusks, 128
mayfly ontogeny, 128
nymphal development, 128
Ephydridae,
Egeria densa, 603
Eucoilinae, 603
Hydrellia, 603
Hydrelliaeucoila egeria, 603
Eriophyidae,
Acaralox, 12
Acari, 12
Aculodes, 12
Poaceae, 12
Estrogen receptor-related receptor (ERR),
molecular cloning, 295
Polyrhachis vicina, 295
5′-/3′-rapid amplification of cDNA ends (RACE), 295
real-time quantitative polymerase chain reaction (PCR), 295
ethnobiology,
cucurbit domestication, 60
geometric morphometrics, 60
range expansion, 60
squash bee, 60
Eucoilinae,
Egeria densa, 603
Ephydridae, 603
Hydrellia, 603
Hydrelliaeucoila egeria, 603
Euplectitae,
Pselaphinae, 406
rove beetles, 406
short-winged mold beetles, 406
Staphylinidae, 406
excavating behavior,
Coptotermes formosanus, 333
mouthparts, 333
tunnel architecture, 333
excavation,
idiothetic, 158
path integration, 158
stigmergy, 158
tunnel, 158
experimental evolution,
adaptation, 181
artificial selection, 181
host shift, 181
oviposition, 181
extended development,
aestivation, 456
galling aphid, 456
life history evolution, 456
Pinaceae, 456

F
fecundity, 413, 476
Asian citrus psyllid, 476
body color, 476
fitness, 413
longevity, 413
pigment, 476
reproductive output, 476
secondary symbionts, 413
survival, 413
fertility,
codling moth, 233
life table parameters, 233
reproduction, 233
Weibull distribution, 233
filter chamber,
alimentary canal, 650
huanglongbing, 650
Liberibacter, 650
Psyllidae, 650
fine motor skills,
functional morphology, 713
grasping, 713
handedness, 713

fire ant,
parasitoid, 937
phorid, 937
Pseudacteon, 937
Solenopsis, 937
fitness,
fecundity, 413
longevity, 413
secondary symbionts, 413
survival, 413
flea beetle,
Altica litigata, 386
herbivore, 386
internal transcribed spacer, 386
mitochondrial DNA, 386
flight behavior,
competitive displacement, 426
Scolytidae, 426
Scolytus multistriatus, 426
Scolytus schevyrewi, 426
flight performance,
food supply, 727
heat stress, 727
olive fruit fly, 727
flightless,
COI, 826
dung beetle, 826
molecular phylogeny, 826
flooding,
Cicindela, 462
hypoxia, 462
immersion, 462
tiger beetles, 462
florivory,
Annonaceae, 492
Costa Rica, 492
Drosophilidae, 492
saproflorivory, 492
food supply,
flight performance, 727
heat stress, 727
olive fruit fly, 727
Forelius,
behavior, 702
Monomorium, 702
Pheidole, 702
Tetramorium, 702
Formosan subterranean termite,
individual workload, 906
tunnel excavation, 906
fruit flies, 310, 819
Anastrepha serpentina, 310
antennal morphology, 310
mango, 819
olfaction, 310
protein, 819
pupae per fruit, 819
S. venulosa, 819
sensilla, 310
functional morphology,
fine motor skills, 713
grasping, 713
handedness, 713

G

galling aphid,
aestivation, 456
extended development, 456
life history evolution, 456
Pinaceae, 456
genetic diversity, 539, 666
Adelges tsugae, 539
allozymes, 539
Apis mellifera, 666
commercial populations, 666
microsatellites, 666
mitochondrial DNA, 666
parthenogenic, 539
genetic variation,
amplified fragment length polymorphism, 151
Bt resistance, 151
population structure, 151
Trichoplusia ni, 151
geometric morphometrics, 1, 60, 1013
apple maggot, 1013
biotypes, 1
canonical variates analysis, 1013
cucurbit domestication, 60
Ephedrus persicae, 1
ethnobiology, 60
ovipositor length, 1013
range expansion, 60
squash bee, 60
variability, 1
gland plasticity,
behavior, 137
life stages, 137
propolis, 137
stingless bees, 137
glassy-winged sharpshooter,
degree-days, 249
developmental threshold, 249
host plants, 249
gonadal development,
Nezara viridula, 303
ovaries, 303
reproductive systems, 303
testes, 303
Grand Canyon,
biogeography, 261
elevation, 261
Odonata, 261
species richness, 261
grasping,
fine motor skills, 713
functional morphology, 713
handedness, 713
ground-dwelling ants,
activity schedule, 1151
ant behavior, 1151
diet, 1151
growth rate,
development, 467
hatching size, 467
interpopulation variation, 467
south Florida, 467
Gryllus,
cryptic species, 367
introduced species, 367
sister species, 367
gustation,
gypsy moth, 1116
sensillum, 1116
styloconic, 1116
ultrastructure, 1116
gut symbionts,
Coptotermes, 498
insect gut, 498
Isoptera, 498
Rhinotermitidae, 498
gypsy moth,
gustation, 1116
sensillum, 1116
styloconic, 1116
ultrastructure, 1116

H

Halyomorpha halys, 39, 608
biocontrol, 39
egg parasitoid, 39
host plant, 608
new species, 39
population biology, 608
seasonal dynamics, 608
Trissolcus halyomorphae, 39
handedness,
fine motor skills, 713
functional morphology, 713
grasping, 713
Harmonia axyridis,
Coccinella septempunctata, 419
Coleomegilla maculata, 419
interspecific interactions, 419
introduced species, 419
hatching size,
development, 467
growth rate, 467
interpopulation variation, 467
south Florida, 467
heat stress,
flight performance, 727
food supply, 727
olive fruit fly, 727
heliconiines,
Laparus, 1105
nutritional ecology, 1105
sugar assay, 1105

Helicoverpa armigera,
body color, 76
host plant, 76
larval diet, 76
polymorphism, 76
Helota,
Helotidae, 48
taxonomic revision, 48
thibetana species group, 48
Helotidae, 48, 785
Helota, 48
Metahelotella, 785
new species, 785
taxonomic revision, 48, 785
thibetana species group, 48
herbivore,
Altica litigata, 386
flea beetle, 386
internal transcribed spacer, 386
mitochondrial DNA, 386
Homalodisca spp.,
electrical penetration graph, 847
Pierce's disease, 847
probing, 847
stylet penetration, 847
Honduras,
mesophilous forest, 959
México, 959
Pseudomenotelus, 959
Pseudomenus, 959
horizontal transfer,
Encarsia, 1050
parthenogenesis, 1050
secondary endosymbiont, 1050
host behavior,
immature development, 288
longevity, 288
morphology, 288
weight, 288
host plant, 76, 210, 249, 608
Anisomorpha buprestoides, 210
body color, 76
color forms, 210
degree-days, 249
developmental threshold, 249
ecology, 210
glassy-winged sharpshooter, 249
Halyomorpha halys, 608
Helicoverpa armigera, 76
intraspecific variation, 210
larval diet, 76
polymorphism, 76
population biology, 608
seasonal dynamics, 608
host searching,
dust/frass, 166
ectoparasitoid, 166
semiochemicals, 166
host shift,
adaptation, 181
artificial selection, 181
experimental evolution, 181
oviposition, 181
huanglongbing, 523, 547, 650
acoustic communication, 547
alimentary canal, 650
antennal morphology, 523
Asian citrus psyllid, 547
citrus greening, 523
citrus greening disease, 547
electron microscopy, 523
filter chamber, 650
Liberibacter, 650
Psyllidae, 650
hybridization,
cryptic, 914
karyotype, 914
mitochondrial DNA, 914
nuclear DNA, 914
Hydrellia,
Egeria densa, 603
Ephydridae, 603
Eucoilinae, 603
Hydrelliaeucoila egeria, 603
Hydrelliaeucoila egeria,
Egeria densa, 603
Ephydridae, 603
Eucoilinae, 603
Hydrellia, 603
hydrolysable tannins,
Choristoneura fumiferana, 717
condensed tannins, 717
insect herbivore, 717
spruce budworm, 717
hypoxia,
Cicindela, 462
flooding, 462
immersion, 462
tiger beetles, 462

I

Ichneumonidae,
Diadegma semiclausum, 532
immature stages, 532
Plutella xylostella, 532
preimaginal development, 532
idiobiont,
competition, 172
occupation, 172
ovipositional resource, 172
idiothetic,
excavation, 158
path integration, 158
stigmergy, 158
tunnel, 158
immature development,
host behavior, 288
longevity, 288
morphology, 288
weight, 288
immature stages,
Diadegma semiclausum, 532
Ichneumonidae, 532
Plutella xylostella, 532
preimaginal development, 532
immersion,
Cicindela, 462
flooding, 462
hypoxia, 462
tiger beetles, 462
Incisitermes,
soldier, 759
Taiwan, 759
taxonomy, 759
winged imago, 759
individual distance,
Coptotermes formosanus, 1158
individual volume, 1158
Reticulitermes flavipes, 1158
self-organized behavior, 1158
individual volume,
Coptotermes formosanus, 1158
individual distance, 1158
Reticulitermes flavipes, 1158
self-organized behavior, 1158
individual workload,
Formosan subterranean termite, 906
tunnel excavation, 906
infochemicals,
egg parasitoids, 1145
Prays oleae, 1145
Y-tube olfactometer, 1145
insect gut,
Coptotermes, 498
gut symbionts, 498
Isoptera, 498
Rhinotermitidae, 498
insect herbivore,
Choristoneura fumiferana, 717
condensed tannins, 717
hydrolysable tannins, 717
spruce budworm, 717
interference competition,
aggression, 201
coexistence, 201
competitor avoidance, 201
dominance hierarchy, 201
internal transcribed spacer,
Altica litigata, 386
flea beetle, 386
herbivore, 386
mitochondrial DNA, 386
interpopulation variation,
development, 467
growth rate, 467
hatching size, 467
south Florida, 467
interspecific interactions,
Coccinella septempunctata, 419
Coleomegilla maculata, 419
Harmonia axyridis, 419

introduced species, 419
intraspecific variation, 210, 868
Anisomorpha buprestoides, 210
color forms, 210
DNA sequence, 868
ecology, 210
host plants, 210
Missouri, 868
Reticulitermes, 868
subterranean termites, 868
introduced species, 367, 419
Coccinella septempunctata, 419
Coleomegilla maculata, 419
cryptic species, 367
Gryllus, 367
Harmonia axyridis, 419
interspecific interactions, 419
sister species, 367
invasive species, 144, 1135
Aedes albopictus, 144, 1135
life history, 1135
mitochondrial DNA gene, 144
nicotinamide adenine dinucleotide dehydrogenase subunit 5, 144
population genetics, 144
population structure, 1135
Isoptera,
Coptotermes, 498
gut symbionts, 498
insect gut, 498
Rhinotermitidae, 498
Israel,
Diptera, 396
Empididae, 396
new species, 396
Rhamphomyia, 396

J

jackknife,
Brevicoryne brassicae, 450
demography, 450
life table, 450
oilseed rape, 450
Jiulongshan Formation, 20, 588
biogeography, 20
China, 588
Middle Jurassic, 20, 588
new genera, 588
palaeoentomology, 20
Pelecinidae, 20
Xyelotomidae, 588
Jurassic,
China, 964
Daohugou, 964
Kalligrammatidae, 964
Neuroptera, 964

K

Kalligrammatidae,
China, 964
Daohugou, 964
Jurassic, 964
Neuroptera, 964
karyotype,
cryptic, 914
hybridization, 914
mitochondrial DNA, 914
nuclear DNA, 914

L

laboratory rearing,
Corimelaena incognita, 1068
descriptions, 1068
life history, 1068
Thyreocoridae, 1068
Laparus,
heliconiines, 1105
nutritional ecology, 1105
sugar assay, 1105
larval diet,
body color, 76
Helicoverpa armigera, 76
host plant, 76
polymorphism, 76
larval morphology,
Capparaceae, 766
Costa Rica, 766
Pyraloidea, 766
systematics, 766
leafhoppers,
biological control, 241
egg age, 241
egg parasitoids, 241
egg structure, 241
Liberibacter,
alimentary canal, 650
filter chamber, 650
huanglongbing, 650
Psyllidae, 650
life cycle, 96, 104, 642
adventive species, 104
Auchenorrhyncha, 642
biological control, 96
development, 642
Diomus terminatus, 96
Melanaphis sacchari, 96
morphology, 96
Pipturus spp., 104
purposely introduced species, 104
reproduction, 642
Xylella fastidiosa, 642
life history, 508, 1068, 1099, 1135
Aedes albopictus, 1135
Argopistes tsekooni, 508
biological control, 508
Corimelaena incognita, 1068
descriptions, 1068
invasive species, 1135
laboratory rearing, 1068
Ligustrum sinense, 508
phasmids, 1099
population structure, 1135
regeneration, 1099
stick insects, 1099
Thyreocoridae, 1068
trade-offs, 1099
life history evolution,
aestivation, 456
extended development, 456
galling aphid, 456
Pinaceae, 456
life stages,
behavior, 137
gland plasticity, 137
propolis, 137
stingless bees, 137
life table,
Brevicoryne brassicae, 450
demography, 450
jackknife, 450
oilseed rape, 450
life table parameters,
codling moth, 233
fertility, 233
reproduction, 233
Weibull distribution, 233
Ligustrum sinense,
Argopistes tsekooni, 508
biological control, 508
life history, 508
longevity, 70, 288, 413
Bactrocera (Dacus) oleae, 70
egg production, 70
fecundity, 413
fitness, 413
host behavior, 288
immature development, 288
morphology, 288
ovarian maturation, 70
relative humidity, 70
secondary symbionts, 413
survival, 413
weight, 288

M

Machaerotinae,
Cicadomorpha, 739
drawing, 739
new species, 739
taxonomy, 739
Macrotermes carbonarius,
caste, 1091
development, 1091
Macrotermes gilvus, 1091
sexual dimorphism, 1091
Macrotermes gilvus,
caste, 1091
development, 1091
Macrotermes carbonarius, 1091
sexual dimorphism, 1091

male feeding,
polygyny, 629
spermatophore color, 629
spermatophore weight, 629
tobacco budworm, 629
mandibular tusks,
abdominal gills, 128
Ephoron leukon, 128
mayfly ontogeny, 128
nymphal development, 128
mango,
fruit flies, 819
protein, 819
pupae per fruit, 819
S. venulosa, 819
mantid,
antipredator strategy, 517
behavior, 517
crypsis, 517
wind, 517
Mantidae,
Masiphya, 842
parasite, 842
Stagmomantis, 842
Tachinidae, 842
Masiphya,
Mantidae, 842
parasite, 842
Stagmomantis, 842
Tachinidae, 842
mass trapping,
active galleries, 809
olive yield, 809
sticky UV-light–pheromone trap, 809
Zeuzera pyrina, 809
mate choice,
Diabrotica virgifera virgifera, 326
mating behavior, 326
reproductive behavior, 326
maternal effects,
compensatory growth, 881
early embryonic mortality, 881
embryogenesis, 881
queen age, 881
mating behavior, 317, 326
Cerambycidae, 317
contact pheromone, 317
cuticular hydrocarbon, 317
Diabrotica virgifera virgifera, 326
mate choice, 326
Monochamus galloprovincialis, 317
reproductive behavior, 326
mating signal,
cryptic species, 747
song species, 747
substrate vibration, 747
systematics, 747
mating system,
copulation duration, 694
Odonata, 694
mayfly ontogeny,
abdominal gills, 128
Ephoron leukon, 128
mandibular tusks, 128
nymphal development, 128
Melanaphis sacchari,
biological control, 96
Diomus terminatus, 96
life cycle, 96
morphology, 96
Meligethinae,
Asteraceae, 998
Nitidulidae, 998
systematics, 998
Tarchonantheae, 998
memory,
biological invasions, 1109
cognition, 1109
mesophilous forest,
Honduras, 959
México, 959
Pseudomenotelus, 959
Pseudomenus, 959
Metahelotella,
Helotidae, 785
new species, 785
taxonomic revision, 785
México,
Honduras, 959
mesophilous forest, 959
Pseudomenotelus, 959
Pseudomenus, 959
microsatellites, 666, 674, 895
Anoplophora glabripennis, 895
Apis mellifera, 666
Asia, 895
Asian longhorned beetle, 895
commercial populations, 666
genetic diversity, 666
mitochondrial DNA, 666, 895
population genetic structure, 674
Tetranychus cinnabarinus, 674
Tetranychus urticae, 674
Middle Jurassic, 20, 588
biogeography, 20
China, 588
Jiulongshan Formation, 20, 588
new genera, 588
palaeoentomology, 20
Pelecinidae, 20
Xyelotomidae, 588
midgut,
starvation, 119
ultrastructure, 119
zoophytophagous, 119
Mindarus abietinus,
discriminant functions, 970
DNA barcodes, 970
Missouri,
DNA sequence, 868
intraspecific variation, 868
Reticulitermes, 868
subterranean termites, 868
mitochondrial COII,
12S rRNA, 684
16S rRNA, 684
biogeography, 684
molecular phylogeny, 684
mitochondrial DNA, 347, 381, 386, 666, 895, 914, 982, 1077
allozyme, 381
Altica litigata, 386
Anoplophora glabripennis, 895
Apis mellifera, 666
Asia, 895, 1077
Asian longhorned beetle, 895
commercial populations, 666
Coptotermes, 1077
cryptic, 914
Dendroctonus pseudotsugae, 982
DNA barcoding, 347
flea beetle, 386
genetic diversity, 666
herbivore, 386
hybridization, 914
internal transcribed spacer, 386
karyotype, 914
microsatellites, 666, 895
morphology, 381
nuclear DNA, 914
Phlebotominae, 347
Phlebotomus chabaudi, 347
Phlebotomus riouxi, 347
phylogenetics, 1077
phytophagous insect, 381
Pseudotsuga, 982
subspecies, 982
subterranean termites, 1077
systematics, 381
mitochondrial DNA gene,
Aedes albopictus, 144
invasive species, 144
nicotinamide adenine dinucleotide dehydrogenase subunit 5, 144
population genetics, 144
molecular cloning,
estrogen receptor-related receptor (ERR), 295
Polyrhachis vicina, 295
5′-/3′-rapid amplification of cDNA ends (RACE), 295
real-time quantitative polymerase chain reaction (PCR), 295

molecular phylogeny, 684, 826
12S rRNA, 684
16S rRNA, 684
biogeography, 684
COI, 826
dung beetle, 826
flightless, 826
mitochondrial COII, 684
Monochamus galloprovincialis,
Cerambycidae, 317
contact pheromone, 317
cuticular hydrocarbon, 317
mating behavior, 317
Monomorium,
behavior, 702
Forelius, 702
Pheidole, 702
Tetramorium, 702
morphology, 96, 288, 381, 556
allozyme, 381
biological control, 96
Diomus terminatus, 96
host behavior, 288
immature development, 288
life cycle, 96
longevity, 288
Melanaphis sacchari, 96
mitochondrial DNA, 381
oviposition, 556
phytophagous insect, 381
stigma, 556
style, 556
systematics, 381
weight, 288
mouthparts,
Coptotermes formosanus, 333
excavating behavior, 333
tunnel architecture, 333
Muscidae,
Callainireinwardtia woodi, 354
Neotropical, 354
new genus, 354
Reinwardtiini, 354
Myzus persicae,
demographic parameters, 437
development, 437
Dichochrysa prasina, 437
prey availability, 437

N

native bee conservation,
pan trapping, 275
Neobellieria,
Diptera, 567
Sarcophaga, 567
systematics, 567
Tolucamyia, 567
Neotropical,
Callainireinwardtia woodi, 354
Muscidae, 354
new genus, 354
Reinwardtiini, 354
Neuroptera,
China, 964
Daohugou, 964
Jurassic, 964
Kalligrammatidae, 964
new genera,
China, 588
Jiulongshan Formation, 588
Middle Jurassic, 588
Xyelotomidae, 588
new genus,
Callainireinwardtia woodi, 354
Muscidae, 354
Neotropical, 354
Reinwardtiini, 354
new species, 39, 396, 739, 785, 1037
biocontrol, 39
cave, 1037
Cicadomorpha, 739
Collembola, 1037
Diptera, 396
drawing, 739
egg parasitoid, 39
Empididae, 396
Halyomorpha halys, 39
Helotidae, 785
Israel, 396
Machaerotinae, 739
Metahelotella, 785
Onychiurus, 1037
Rhamphomyia, 396
taxonomic revision, 785
taxonomy, 739
Trissolcus halyomorphae, 39
Nezara viridula,
gonadal development, 303
ovaries, 303
reproductive systems, 303
testes, 303
nicotinamide adenine dinucleotide dehydrogenase subunit 5,
Aedes albopictus, 144
invasive species, 144
mitochondrial DNA gene, 144
population genetics, 144
Nitidulidae,
Asteraceae, 998
Meligethinae, 998
systematics, 998
Tarchonantheae, 998
nuclear DNA,
cryptic, 914
hybridization, 914
karyotype, 914
mitochondrial DNA, 914
nutritional ecology,
heliconiines, 1105
Laparus, 1105
sugar assay, 1105
nymphal development,
abdominal gills, 128
Ephoron leukon, 128
mandibular tusks, 128
mayfly ontogeny, 128

O

occupation,
competition, 172
idiobiont, 172
ovipositional resource, 172
Odonata, 261, 694
biogeography, 261
copulation duration, 694
elevation, 261
Grand Canyon, 261
mating system, 694
species richness, 261
Oebalus pugnax,
color changes, 638
development, 638
rice stink bug eggs, 638
temperature, 638
oilseed rape,
Brevicoryne brassicae, 450
demography, 450
jackknife, 450
life table, 450
okra-leaf,
cotton, 196
Creontiades signatus, 196
oviposition, 196
trichomes, 196
olfaction,
Anastrepha serpentina, 310
antennal morphology, 310
fruit flies, 310
sensilla, 310
olive fruit fly,
flight performance, 727
food supply, 727
heat stress, 727
olive yield,
active galleries, 809
mass trapping, 809
sticky UV-light–pheromone trap, 809
Zeuzera pyrina, 809
Onychiurus,
cave, 1037
Collembola, 1037
new species, 1037
oocyte,
alates, 889
reproduction, 889
sperm energy source, 889
spermatheca, 889
Opuntia ficus-indica,
adult body weights, 445
egg incubation, 445
egg production, 445

Orthoptera,
acridids, 339
Caelifera, 339
catantopids, 339
pygmy grasshoppers, 339
ovarian maturation,
Bactrocera (*Dacus*) *oleae*, 70
egg production, 70
longevity, 70
relative humidity, 70
ovaries,
gonadal development, 303
Nezara viridula, 303
reproductive systems, 303
testes, 303
oviposition, 181, 196, 556
adaptation, 181
artificial selection, 181
cotton, 196
Creontiades signatus, 196
experimental evolution, 181
host shift, 181
morphology, 556
okra-leaf, 196
stigma, 556
style, 556
trichomes, 196
ovipositional resource,
competition, 172
idiobiont, 172
occupation, 172
ovipositor length,
apple maggot, 1013
canonical variates analysis, 1013
geometric morphometrics, 1013

P

Pachnaeus litus,
biological control, 112
Diaprepes abbreviatus, 112
egg parasitoid, 112
palaeoentomology,
biogeography, 20
Jiulongshan Formation, 20
Middle Jurassic, 20
Pelecinidae, 20
pan trapping,
native bee conservation, 275
parasite,
Mantidae, 842
Masiphya, 842
Stagmomantis, 842
Tachinidae, 842
parasitoid, 925, 937
Binodoxys communis, 925
Binodoxys koreanus, 925
cryptic species, 925
fire ant, 937
phorid, 937
Pseudacteon, 937
Solenopsis, 937
soybean aphid, 925
parthenogenesis,
Encarsia, 1050
horizontal transfer, 1050
secondary endosymbiont, 1050
parthenogenic,
Adelges tsugae, 539
allozymes, 539
genetic diversity, 539
path integration,
excavation, 158
idiothetic, 158
stigmergy, 158
tunnel, 158
Pelecinidae,
biogeography, 20
Jiulongshan Formation, 20
Middle Jurassic, 20
palaeoentomology, 20
Pezothrips kellyanus,
citrus, 835
developmental time, 835
temperature thresholds, 835
thermal requirements, 835
phasmids,
life history, 1099
regeneration, 1099
stick insects, 1099
trade-offs, 1099
Pheidole,
behavior, 702
Forelius, 702
Monomorium, 702
Tetramorium, 702
Phlebotominae,
DNA barcoding, 347
mitochondrial DNA, 347
Phlebotomus chabaudi, 347
Phlebotomus riouxi, 347
Phlebotomus chabaudi,
DNA barcoding, 347
mitochondrial DNA, 347
Phlebotominae, 347
Phlebotomus riouxi, 347
Phlebotomus riouxi,
DNA barcoding, 347
mitochondrial DNA, 347
Phlebotominae, 347
Phlebotomus chabaudi, 347
phorid,
fire ant, 937
parasitoid, 937
Pseudacteon, 937
Solenopsis, 937
phylogenetics,
Asia, 1077
Coptotermes, 1077
mitochondrial DNA, 1077
subterranean termites, 1077
phylogeny,
Aubehydrini, 797
chaetotaxy, 797
diving beetles, 797
Dytiscidae, 797
phytophagous insect,
allozyme, 381
mitochondrial DNA, 381
morphology, 381
systematics, 381
Phytoseiidae,
predacious mites, 617
satsuma mandarin, 617
Stigmaeidae, 617
Typhlodromalus peregrinus, 617
Pierce's disease,
electrical penetration graph, 847
Homalodisca spp., 847
probing, 847
stylet penetration, 847
pigment,
Asian citrus psyllid, 476
body color, 476
fecundity, 476
reproductive output, 476
Pinaceae,
aestivation, 456
extended development, 456
galling aphid, 456
life history evolution, 456
Pipturus spp.,
adventive species, 104
life cycle, 104
purposely introduced species, 104
Plutella xylostella,
Diadegma semiclausum, 532
Ichneumonidae, 532
immature stages, 532
preimaginal development, 532
Poaceae,
Acaralox, 12
Acari, 12
Aculodes, 12
Eriophyidae, 12
Podapolipidae,
Acari, 1062
Carabidae, 1062
taxonomy, 1062
polygyny,
male feeding, 629
spermatophore color, 629
spermatophore weight, 629
tobacco budworm, 629
polymerase chain reaction-restriction fragment length polymorphism,
Reticulitermes, 1029
16S rRNA, 1029
survey, 1029

termite, 1029
polymorphism,
body color, 76
Helicoverpa armigera, 76
host plant, 76
larval diet, 76
Polyrhachis vicina,
estrogen receptor-related receptor (ERR), 295
molecular cloning, 295
5′-/3′-rapid amplification of cDNA ends (RACE), 295
real-time quantitative polymerase chain reaction (PCR), 295
population biology,
Halyomorpha halys, 608
host plant, 608
seasonal dynamics, 608
population genetic structure,
microsatellites, 674
Tetranychus cinnabarinus, 674
Tetranychus urticae, 674
population genetics,
Aedes albopictus, 144
invasive species, 144
mitochondrial DNA gene, 144
nicotinamide adenine dinucleotide dehydrogenase subunit 5, 144
population structure, 151, 1135
Aedes albopictus, 1135
amplified fragment length polymorphism, 151
Bt resistance, 151
genetic variation, 151
invasive species, 1135
life history, 1135
Trichoplusia ni, 151
powdery mildew,
biological control, 484
Psyllobora vigintimaculata, 484
Prays oleae,
egg parasitoids, 1145
infochemicals, 1145
Y-tube olfactometer, 1145
predacious mites,
Phytoseiidae, 617
satsuma mandarin, 617
Stigmaeidae, 617
Typhlodromalus peregrinus, 617
preimaginal development,
Diadegma semiclausum, 532
Ichneumonidae, 532
immature stages, 532
Plutella xylostella, 532
prey availability,
demographic parameters, 437
development, 437
Dichochrysa prasina, 437
Myzus persicae, 437
probing,
electrical penetration graph, 847
Homalodisca spp., 847
Pierce's disease, 847
stylet penetration, 847
propolis,
behavior, 137
gland plasticity, 137
life stages, 137
stingless bees, 137
protein,
fruit flies, 819
mango, 819
pupae per fruit, 819
S. venulosa, 819
Pselaphinae,
Euplectitae, 406
rove beetles, 406
short-winged mold beetles, 406
Staphylinidae, 406
Pseudacteon,
fire ant, 937
parasitoid, 937
phorid, 937
Solenopsis, 937
Pseudomenotelus,
Honduras, 959
mesophilous forest, 959
México, 959
Pseudomenus, 959
Pseudomenus,
Honduras, 959
mesophilous forest, 959
México, 959
Pseudomenotelus, 959
Pseudotsuga,
Dendroctonus pseudotsugae, 982
mitochondrial DNA, 982
subspecies, 982
Psyllidae,
alimentary canal, 650
filter chamber, 650
huanglongbing, 650
Liberibacter, 650
Psyllobora vigintimaculata,
biological control, 484
powdery mildew, 484
pupae per fruit,
fruit flies, 819
mango, 819
protein, 819
S. venulosa, 819
purposely introduced species,
adventive species, 104
life cycle, 104
Pipturus spp., 104
pygmy grasshoppers,
acridids, 339
Caelifera, 339
catantopids, 339
Orthoptera, 339
Pyraloidea,
Capparaceae, 766
Costa Rica, 766
larval morphology, 766
systematics, 766

Q

queen age,
compensatory growth, 881
early embryonic mortality, 881
embryogenesis, 881
maternal effects, 881

R

range expansion,
cucurbit domestication, 60
ethnobiology, 60
geometric morphometrics, 60
squash bee, 60
5′-/3′-rapid amplification of cDNA ends (RACE),
estrogen receptor-related receptor (ERR), 295
molecular cloning, 295
Polyrhachis vicina, 295
real-time quantitative polymerase chain reaction (PCR), 295
real-time quantitative polymerase chain reaction (PCR),
estrogen receptor-related receptor (ERR), 295
molecular cloning, 295
Polyrhachis vicina, 295
5′-/3′-rapid amplification of cDNA ends (RACE), 295
regeneration,
life history, 1099
phasmids, 1099
stick insects, 1099
trade-offs, 1099
Reinwardtiini,
Callainireinwardtia woodi, 354
Muscidae, 354
Neotropical, 354
new genus, 354
relative humidity,
Bactrocera (*Dacus*) *oleae*, 70
egg production, 70
longevity, 70
ovarian maturation, 70
reproduction, 85, 233, 642, 889
alates, 889
Auchenorrhyncha, 642
codling moth, 233
development, 642

fertility, 233
life cycle, 642
life table parameters, 233
oocyte, 889
sperm energy source, 889
spermatheca, 889
sterile insect technique, 85
symbiosis, 85
Weibull distribution, 233
Xylella fastidiosa, 642
reproductive behavior,
Diabrotica virgifera virgifera, 326
mate choice, 326
mating behavior, 326
reproductive isolation,
biodiversity, 360
social spider mites, 360
Tetranychidae, 360
reproductive output,
Asian citrus psyllid, 476
body color, 476
fecundity, 476
pigment, 476
reproductive systems,
gonadal development, 303
Nezara viridula, 303
ovaries, 303
testes, 303
Reticulitermes, 868, 1029
DNA sequence, 868
intraspecific variation, 868
Missouri, 868
polymerase chain reaction-restriction fragment length polymorphism, 1029
16S rRNA, 1029
subterranean termites, 868
survey, 1029
termite, 1029
Reticulitermes flavipes,
Coptotermes formosanus, 1158
individual distance, 1158
individual volume, 1158
self-organized behavior, 1158
Rhamphomyia,
Diptera, 396
Empididae, 396
Israel, 396
new species, 396
Rhinotermitidae,
Coptotermes, 498
gut symbionts, 498
insect gut, 498
Isoptera, 498
rice stink bug eggs,
color changes, 638
development, 638
Oebalus pugnax, 638
temperature, 638
rove beetles, 189, 406
12s, 189
16s, 189
Aleocharinae, 189
Euplectitae, 406
Pselaphinae, 406
short-winged mold beetles, 406
Staphylinidae, 189, 406

S

12s,
16s, 189
Aleocharinae, 189
rove beetles, 189
Staphylinidae, 189
12S rRNA,
16S rRNA, 684
biogeography, 684
mitochondrial COII, 684
molecular phylogeny, 684
16s,
12s, 189
Aleocharinae, 189
rove beetles, 189
Staphylinidae, 189
16S rRNA, 684, 1029
12S rRNA, 684
biogeography, 684
mitochondrial COII, 684
molecular phylogeny, 684
polymerase chain reaction-restriction fragment length polymorphism, 1029
Reticulitermes, 1029
survey, 1029
termite, 1029
S. venulosa,
fruit flies, 819
mango, 819
protein, 819
pupae per fruit, 819
saproflorivory,
Annonaceae, 492
Costa Rica, 492
Drosophilidae, 492
florivory, 492
Sarcophaga,
Diptera, 567
Neobellieria, 567
systematics, 567
Tolucamyia, 567
satsuma mandarin,
Phytoseiidae, 617
predacious mites, 617
Stigmaeidae, 617
Typhlodromalus peregrinus, 617
Scolytidae,
competitive displacement, 426
flight behavior, 426
Scolytus multistriatus, 426
Scolytus schevyrewi, 426
Scolytinae,
Scolytus abietis, 597
Scolytus opacus, 597
synonymy, 597
Scolytus abietis,
Scolytinae, 597
Scolytus opacus, 597
synonymy, 597
Scolytus multistriatus,
competitive displacement, 426
flight behavior, 426
Scolytidae, 426
Scolytus schevyrewi, 426
Scolytus opacus,
Scolytinae, 597
Scolytus abietis, 597
synonymy, 597
Scolytus schevyrewi,
competitive displacement, 426
flight behavior, 426
Scolytidae, 426
Scolytus multistriatus, 426
seasonal dynamics,
Halyomorpha halys, 608
host plant, 608
population biology, 608
secondary endosymbiont,
Encarsia, 1050
horizontal transfer, 1050
parthenogenesis, 1050
secondary symbionts,
fecundity, 413
fitness, 413
longevity, 413
survival, 413
self-organized behavior,
Coptotermes formosanus, 1158
individual distance, 1158
individual volume, 1158
Reticulitermes flavipes, 1158
semiochemicals,
dust/frass, 166
ectoparasitoid, 166
host searching, 166
sensilla,
Anastrepha serpentina, 310
antennal morphology, 310
fruit flies, 310
olfaction, 310
sensillum,
gustation, 1116
gypsy moth, 1116
styloconic, 1116
ultrastructure, 1116
sexual dimorphism,
caste, 1091
development, 1091
Macrotermes carbonarius, 1091
Macrotermes gilvus, 1091

short-winged mold beetles,
Euplectitae, 406
Pselaphinae, 406
rove beetles, 406
Staphylinidae, 406
sister species,
cryptic species, 367
Gryllus, 367
introduced species, 367
social spider mites,
biodiversity, 360
reproductive isolation, 360
Tetranychidae, 360
soldier,
Incisitermes, 759
Taiwan, 759
taxonomy, 759
winged imago, 759
Solenopsis,
fire ant, 937
parasitoid, 937
phorid, 937
Pseudacteon, 937
song species,
cryptic species, 747
mating signal, 747
substrate vibration, 747
systematics, 747
south Florida,
development, 467
growth rate, 467
hatching size, 467
interpopulation variation, 467
soybean aphid,
Binodoxys communis, 925
Binodoxys koreanus, 925
cryptic species, 925
parasitoids, 925
species richness,
biogeography, 261
elevation, 261
Grand Canyon, 261
Odonata, 261
sperm energy source,
alates, 889
oocyte, 889
reproduction, 889
spermatheca, 889
spermatheca,
alates, 889
oocyte, 889
reproduction, 889
sperm energy source, 889
spermatophore color,
male feeding, 629
polygyny, 629
spermatophore weight, 629
tobacco budworm, 629
spermatophore weight,
male feeding, 629
polygyny, 629
spermatophore color, 629
tobacco budworm, 629
spruce budworm,
Choristoneura fumiferana, 717
condensed tannins, 717
hydrolysable tannins, 717
insect herbivore, 717
squash bee,
cucurbit domestication, 60
ethnobiology, 60
geometric morphometrics, 60
range expansion, 60
Stagmomantis,
Mantidae, 842
Masiphya, 842
parasite, 842
Tachinidae, 842
Staphylinidae, 189, 406
12s, 189
16s, 189
Aleocharinae, 189
Euplectitae, 406
Pselaphinae, 406
rove beetles, 189, 406
short-winged mold beetles, 406
starvation,
midgut, 119
ultrastructure, 119
zoophytophagous, 119
sterile insect technique,
reproduction, 85
symbiosis, 85
stick insects,
life history, 1099
phasmids, 1099
regeneration, 1099
trade-offs, 1099
sticky UV-light–pheromone trap,
active galleries, 809
mass trapping, 809
olive yield, 809
sticky UV-light–pheromone trap, 809
Zeuzera pyrina, 809
stigma,
morphology, 556
oviposition, 556
style, 556
Stigmaeidae,
Phytoseiidae, 617
predacious mites, 617
satsuma mandarin, 617
Typhlodromalus peregrinus, 617
stigmergy,
excavation, 158
idiothetic, 158
path integration, 158
tunnel, 158
stingless bees,
behavior, 137
gland plasticity, 137
life stages, 137
propolis, 137
style,
morphology, 556
oviposition, 556
stigma, 556
stylet penetration,
electrical penetration graph, 847
Homalodisca spp., 847
Pierce's disease, 847
probing, 847
styloconic,
gustation, 1116
gypsy moth, 1116
sensillum, 1116
ultrastructure, 1116
subspecies,
Dendroctonus pseudotsugae, 982
mitochondrial DNA, 982
Pseudotsuga, 982
substrate vibration,
cryptic species, 747
mating signal, 747
song species, 747
systematics, 747
subterranean termites, 868, 1077
Asia, 1077
Coptotermes, 1077
DNA sequence, 868
intraspecific variation, 868
Missouri, 868
mitochondrial DNA, 1077
phylogenetics, 1077
Reticulitermes, 868
sugar assay,
heliconiines, 1105
Laparus, 1105
nutritional ecology, 1105
survey,
polymerase chain reaction-restriction fragment length polymorphism, 1029
Reticulitermes, 1029
16S rRNA, 1029
termite, 1029
survival,
fecundity, 413
fitness, 413
longevity, 413
secondary symbionts, 413
symbiosis,
reproduction, 85
sterile insect technique, 85
synonymy,
Scolytinae, 597
Scolytus abietis, 597
Scolytus opacus, 597

systematics, 381, 567, 747, 766, 998
allozyme, 381
Asteraceae, 998
Capparaceae, 766
Costa Rica, 766
cryptic species, 747
Diptera, 567
larval morphology, 766
mating signal, 747
Meligethinae, 998
mitochondrial DNA, 381
morphology, 381
Neobellieria, 567
Nitidulidae, 998
phytophagous insect, 381
Pyraloidea, 766
Sarcophaga, 567
song species, 747
substrate vibration, 747
Tarchonantheae, 998
Tolucamyia, 567

T

Tachinidae,
Mantidae, 842
Masiphya, 842
parasite, 842
Stagmomantis, 842
Taiwan,
Incisitermes, 759
soldier, 759
taxonomy, 759
winged imago, 759
Tarchonantheae,
Asteraceae, 998
Meligethinae, 998
Nitidulidae, 998
systematics, 998
taxonomic revision, 48, 785
Helota, 48
Helotidae, 48, 785
Metahelotella, 785
new species, 785
thibetana species group, 48
taxonomy, 739, 759, 1062
Acari, 1062
Carabidae, 1062
Cicadomorpha, 739
drawing, 739
Incisitermes, 759
Machaerotinae, 739
new species, 739
Podapolipidae, 1062
soldier, 759
Taiwan, 759
winged imago, 759
temperature,
color changes, 638
development, 638
Oebalus pugnax, 638
rice stink bug eggs, 638
temperature thresholds,
citrus, 835
developmental time, 835
Pezothrips kellyanus, 835
thermal requirements, 835
termite,
polymerase chain reaction-restriction fragment length polymorphism, 1029
Reticulitermes, 1029
16S rRNA, 1029
survey, 1029
testes,
gonadal development, 303
Nezara viridula, 303
ovaries, 303
reproductive systems, 303
Tetramorium,
behavior, 702
Forelius, 702
Monomorium, 702
Pheidole, 702
Tetranychidae,
biodiversity, 360
reproductive isolation, 360
social spider mites, 360
Tetranychus cinnabarinus,
microsatellites, 674
population genetic structure, 674
Tetranychus urticae, 674
Tetranychus urticae,
microsatellites, 674
population genetic structure, 674
Tetranychus cinnabarinus, 674
thermal requirements,
citrus, 835
developmental time, 835
Pezothrips kellyanus, 835
temperature thresholds, 835
thibetana species group,
Helota, 48
Helotidae, 48
taxonomic revision, 48
Thyreocoridae,
Corimelaena incognita, 1068
descriptions, 1068
laboratory rearing, 1068
life history, 1068
tiger beetles,
Cicindela, 462
flooding, 462
hypoxia, 462
immersion, 462
tobacco budworm,
male feeding, 629
polygyny, 629
spermatophore color, 629
spermatophore weight, 629
Tolucamyia,
Diptera, 567
Neobellieria, 567
Sarcophaga, 567
systematics, 567
trade-offs,
life history, 1099
phasmids, 1099
regeneration, 1099
stick insects, 1099
trichomes,
cotton, 196
Creontiades signatus, 196
okra-leaf, 196
oviposition, 196
Trichoplusia ni,
amplified fragment length polymorphism, 151
Bt resistance, 151
genetic variation, 151
population structure, 151
trimethylpyrazine,
dihydropyrazine, 560
2,5-dimethylpyrazine, 560
Trissolcus halyomorphae,
biocontrol, 39
egg parasitoid, 39
Halyomorpha halys, 39
new species, 39
tunnel,
excavation, 158
idiothetic, 158
path integration, 158
stigmergy, 158
tunnel architecture,
Coptotermes formosanus, 333
excavating behavior, 333
mouthparts, 333
tunnel excavation,
Formosan subterranean termite, 906
individual workload, 906
Typhlodromalus peregrinus,
Phytoseiidae, 617
predacious mites, 617
satsuma mandarin, 617
Stigmaeidae, 617

U

ultrastructure, 119, 1116
gustation, 1116
gypsy moth, 1116
midgut, 119
sensillum, 1116
starvation, 119
styloconic, 1116
zoophytophagous, 119

V

variability,
biotypes, 1
Ephedrus persicae, 1

geometric morphometrics, 1

W

Weibull distribution,
codling moth, 233
fertility, 233
life table parameters, 233
reproduction, 233
weight,
host behavior, 288
immature development, 288
longevity, 288
morphology, 288
wind,
antipredator strategy, 517
behavior, 517
crypsis, 517
mantid, 517
winged imago,
Incisitermes, 759
soldier, 759
Taiwan, 759
taxonomy, 759
Xyelotomidae,
China, 588
Jiulongshan Formation, 588
Middle Jurassic, 588
new genera, 588
Xylella fastidiosa,
Auchenorrhyncha, 642
development, 642
life cycle, 642
reproduction, 642

Y

Y-tube olfactometer,
egg parasitoids, 1145
infochemicals, 1145
Prays oleae, 1145

Z

Zeuzera pyrina,
active galleries, 809
mass trapping, 809
olive yield, 809
sticky UV-light–pheromone trap, 809
zoophytophagous,
midgut, 119
starvation, 119
ultrastructure, 119